

# TCL1A Single-Nucleotide Polymorphisms and Estrogen-Mediated Toll-Like Receptor-MYD88-Dependent Nuclear Factor- $\kappa$ B Activation: Single-Nucleotide Polymorphism- and Selective Estrogen Receptor Modulator-Dependent Modification of Inflammation and Immune Response

Ming-Fen Ho, James N. Ingle, Tim Bongartz, Krishna R. Kalari, Paul E. Goss, Lois E. Shepherd, Taisei Mushiroda, Michiaki Kubo, Liewei Wang, and Richard M. Weinshilboum

Division of Clinical Pharmacology, Department of Molecular Pharmacology and Experimental Therapeutics (M.-F.H., L.W., R.M.W.), Division of Medical Oncology, Department of Oncology (J.N.I.), and Division of Biomedical Statistics and Informatics, Department of Health Sciences Research (K.R.K.), Mayo Clinic, Rochester, Minnesota; Department of Emergency Medicine, Vanderbilt Medical Center, Nashville, Tennessee (T.B.); Division of Hematology/Oncology, Department of Medicine, Massachusetts General Hospital Cancer Center, Boston, Massachusetts (P.E.G.); NCIC Clinical Trials Group, Kingston, Ontario Canada (L.E.S.); and RIKEN Center for Integrative Medical Science, Tsurumi-ku, Yokohama, Japan (T.M., M.K.)

Received January 11, 2017; accepted May 30, 2017

## ABSTRACT

In a previous genome-wide association study (GWAS) for musculoskeletal adverse events during aromatase inhibitor therapy for breast cancer, we reported that single nucleotide polymorphisms (SNPs) near the *TCL1A* gene were associated with this adverse drug reaction. Functional genomic studies showed that *TCL1A* expression was induced by estradiol, but only in cells with the variant sequence for the top GWAS SNP (rs11849538), a SNP that created a functional estrogen response element. In addition, *TCL1A* genotype influenced the downstream expression of a series of cytokines and chemokines and had a striking effect on nuclear factor  $\kappa$ B (NF- $\kappa$ B) transcriptional activity. Furthermore, this SNP-dependent regulation could be reversed by selective ER modulators (SERMs). The present study was designed to pursue mechanisms underlying *TCL1A* SNP-mediated, estrogen-dependent NF- $\kappa$ B activation.

Functional genomic studies were performed using a panel of 300 lymphoblastoid cell lines for which we had generated genome-wide SNP and gene expression data. It is known that toll-like receptors (TLRs) can regulate NF- $\kappa$ B signaling by a process that requires the adaptor protein MYD88. We found that TLR2, TLR7, TLR9, and TLR10 expression, as well as that of MYD88, could be modulated by *TCL1A* in a SNP and estrogen-dependent fashion and that these effects were reversed in the presence of SERMs. Furthermore, MYD88 inhibition blocked the *TCL1A* SNP and estrogen-dependent NF- $\kappa$ B activation, as well as protein-protein interaction between *TCL1A* and MYD88. These observations greatly expand the range of pathways influenced by *TCL1A* genotype and raise the possibility that this estrogen- and SNP-dependent regulation might be altered pharmacologically by SERMs.


## Introduction

In a previous genome-wide association study (GWAS) addressing musculoskeletal adverse events in breast cancer patients treated with aromatase inhibitors, we identified single nucleotide polymorphisms (SNPs) 3' of the T-cell leukemia/lymphoma 1A gene (*TCL1A*) that were associated with this adverse drug

reaction (Ingle et al., 2010). The minor allele frequency for the *TCL1A* SNP (rs11849538) is ~19% in European, African, Asian, and Caucasian Americans according to the 1000 Genome Project (Ingle et al., 2010). We subsequently performed a series of functional genomic studies using as a model system lymphoblastoid cell lines (LCLs) for which we had generated dense genome-wide genomic data. Those studies showed that *TCL1A* expression was upregulated by estradiol ( $E_2$ ), but only in cell lines homozygous for the variant *TCL1A* genotype for the top GWAS SNP (rs11849538) (Fig. 1A), a SNP that created a functional estrogen response element (ERE) (Ingle et al., 2010). We subsequently found that there were at least three SNPs 3' of

This work was supported in part by the National Institutes of Health (NIH) National Institute of General Medical Sciences [U19 GM61388] and NIH [P50 CA11620, RO1 CA196648, RO1 GM28157, and RO1 CA138461], and the Breast Cancer Research Foundation.  
<https://doi.org/10.1124/mol.117.108340>

**ABBREVIATIONS:** 4-OH-TAM, 4-hydroxytamoxifen;  $E_2$ , estradiol; ER, estrogen receptor; ERE, estrogen response element; FBS, fetal bovine serum; GWAS, genome-wide association study; ICI, fulvestrant; IP, immunoprecipitation; IRAK, interleukin-1R-activating kinase; LCL, lymphoblastoid cell lines; NF- $\kappa$ B, nuclear factor  $\kappa$ B; PCR, polymerase chain reaction; SERM, selective estrogen receptor modulator; SNP, single nucleotide polymorphisms; *TCL1A*, T-cell leukemia/lymphoma 1A; TLR, toll-like receptor; WT, wild-type.



**Fig. 1.** The SNP- and estrogen-dependent effects on mRNA expression of *TCL1A*, cytokines, and chemokines in LCLs. (A) Schematic diagrams of the two *TCL1A* SNPs, rs7359033 and rs7160302, in tight LD with rs11849538, the top hit signal from the MA.27 GWAS. Locations of EREs are shown as boxes for these three SNPs that map between the 3'-termini of *TCL1A* and *TCL1B*. ER blockade by 4-hydroxytamoxifen (4OH) or fulvestrant (ICI) treatment resulted in the reversal of *TCL1A* SNP and estrogen-dependent *TCL1A* expression patterns (B and C) and downstream effects on the expression of CCR6, CCL20, IL17RA, and IL17A (D–G). Values are mean ± S.E.M. of three assays. \**P* < 0.0001. Adapted from Fig. 2 and Fig. 3 in Ho et al. (2016).

*TCL1A*, rs11849538, rs7359033, and rs7160302 (Fig. 1A)—all in tight linkage disequilibrium (LD)—that appeared to act in concert to influence the E<sub>2</sub>-dependent induction of *TCL1A* expression, in part because SNPs near, but not in, an ERE can affect estrogen receptor (ER) binding to the ERE and subsequent transcription (Ho et al., 2016). In addition, induction of *TCL1A* by E<sub>2</sub> was associated downstream with variation in the expression of a series of proinflammatory cytokines (Liu et al., 2012) and chemokines (Ho et al., 2016), as well as their receptors (Fig. 1, D–G), and with striking variation in nuclear factor  $\kappa$ B (NF- $\kappa$ B) transcriptional activity (Liu et al., 2012). For example, we found that the expression of immune mediators, such as the C-C motif chemokine receptor 6 (*CCR6*) and its only known ligand, *CCL20*, as well as the cytokine receptor *IL17RA* and its ligand, *IL17A*, could be modulated by *TCL1A* in a SNP-estrogen-dependent fashion (Ho et al., 2016). In addition, we found that the SNP and estrogen-dependent induction of *TCL1A* expression could be reversed after ER blockade, that is, when cells were exposed to 4-hydroxytamoxifen (4-OH-TAM), an active metabolite of tamoxifen, or to fulvestrant (formerly *ICI 182,780*) (Fig. 1, B and C), an ER antagonist (Liu et al., 2012; Ho et al., 2016). Specifically, cells homozygous for the wild-type (WT) genotype for the *TCL1A* SNPs displayed increased *TCL1A* expression, whereas cells homozygous for the variant genotype showed decreased expression in the presence of these drugs—the opposite of the situation seen after estrogen exposure alone (Fig. 1, B and C). Even more striking, downstream receptors for inflammatory mediators (*CCR6* and *IL17RA*) responded in a parallel fashion, indicating that *TCL1A* was upstream of the cytokine and chemokine receptors in terms of these pathways (Ho et al., 2016). It should be emphasized that the SNPs involved were those 3' of *TCL1A*, not SNPs that mapped to the *IL17RA* or *CCR6* genes (Ho et al., 2016). Finally, SNP and estrogen-dependent *TCL1A* induction resulted in significantly increased NF- $\kappa$ B transcriptional activity after ER blockade, but only for the variant genotype (Liu et al., 2012). Taken as a whole, these observations suggest that *TCL1A* genotype is capable of influencing the expression of a series of genes that play important roles in inflammation and immunity, and they also raise the possibility of pharmacologic approaches (i.e., exposure to SERMs) that could be used to modify these effects.

Although we have studied the transcriptional regulation of cytokines and chemokines and their receptors in relation to *TCL1A* SNPs and estrogens (Liu et al., 2012; Ho et al., 2016), significant gaps remain in our knowledge, particularly with regard to mechanisms that might be involved in the effect of *TCL1A* on NF- $\kappa$ B signaling. NF- $\kappa$ B activation is a multifactorial, complex process involving many biologic pathways, such as DNA damage (McCool and Miyamoto, 2012), proteasome-mediated degradation (Hayden et al., 2006), signal transduction via stimulation of T cells, or exposure to inflammatory molecules (Takeda et al., 2008). The expression of NF- $\kappa$ B varies among cell types and can be event-specific, depending on the stimulus (Tak and Firestein, 2001). Toll-like receptors (TLRs) are crucial immune mediators and can have a profound effect on inflammation as a result, in part, of NF- $\kappa$ B activation (Tak and Firestein, 2001; Kawasaki and Kawai, 2014). In the present experiments, we set out to study mechanisms that might be involved in the *TCL1A* SNP-mediated, estrogen-dependent activation of NF- $\kappa$ B (Liu et al., 2012). Specifically, we aimed to determine whether the expression of TLRs and their adaptor molecules might be altered in a *TCL1A* SNP-dependent fashion and, if so, their possible contribution to

the *TCL1A* SNP-mediated, estrogen-dependent activation of NF- $\kappa$ B. In addition, we tested the hypothesis that SERMs might be able to modulate these effects. In summary, the experiments described subsequently involved the application of functional genomic studies to identify mechanisms underlying the effect of *TCL1A* SNPs on the regulation of inflammatory mediators.

## Materials and Methods

**Ethics Statement.** The Mayo Clinic Institutional Review Board determined that this work did not require their review or approval.

**Human Variation Panel Lymphoblastoid Cell Lines.** A Human Variation Panel lymphoblastoid cell line (LCL) model system consisting of 300 LCLs from healthy subjects of three ethnicities (100 European American, 100 African American, and 100 Han Chinese American) was used to perform these studies. This cell line model system provides genome-wide mRNA expression as determined by Affymetrix U133 2.0 Plus GeneChip expression array as well as genome-wide SNP data generated with Illumina 550K and 510S SNP BeadChip SNP arrays (Illumina, San Diego, CA). The genotype data were used to impute approximately seven million SNPs per cell line.

**RNA Interference and Transfection.** We purchased siRNA (*TCL1A* and *MYD88*) and negative controls from Dharmacon (Chicago, IL). LCLs were transfected with siRNA by electroporation using the nucleofector kit (Lonza, Allendale, NJ). Briefly, the electroporation reaction contained  $2 \times 10^6$  cells, 100  $\mu$ l of nucleofector solution, and 1  $\mu$ M siRNA. After electroporation, cells were transferred into 12-well plates containing pre-equilibrated RPMI medium.

**Drug Treatment.** LCLs with known genotypes were cultured in RPMI 1640 media (Cellgro, Manassas, VA) supplemented with 15% fetal bovine serum (FBS; Atlanta Biologicals, Flowery Branch, GA). Before estrogen treatment, cells were cultured in RPMI media containing 5% (v/v) charcoal stripped FBS for 24 hours. The cells were then cultured in FBS-free RPMI media for another 24 hours and were treated with 0.1 nM E<sub>2</sub> (Sigma, St. Louis, MO) for 24 hours. In some experiments, cells were treated with  $10^{-7}$   $\mu$ M 4-hydroxytamoxifen (4-OH-TAM) (Sigma) or fulvestrant (formerly *ICI 182,780*) (Sigma), followed by E<sub>2</sub> treatment. *MYD88* inhibitor peptide was purchased from Novus Biologicals (Littleton, CO).

**Real-Time Polymerase Chain Reaction.** RNA was isolated from the cells (Zymo, Irvine, CA). The polymerase chain reaction (PCR) mixture contained 50 ng of total RNA, 5  $\mu$ l of 2 $\times$  SYBR Green real-time PCR mix (Affymetrix, Santa Clara, CA), 0.1  $\mu$ l of DNA polymerase enzyme, 1  $\mu$ l of gene-specific primer (Qiagen, Valencia, CA), and distilled water to achieve 10  $\mu$ l per reaction. Real-time PCR reactions were performed in duplicate using Applied Biosystems ViiA 7 real-time PCR System (Life Technologies, Carlsbad, CA). The 2<sup>- $\Delta\Delta$ Ct</sup> method was used for statistical data analysis.

**Western Blot Analysis.** Protein samples were used to perform electrophoresis, followed by transfer to a polyvinylidene fluoride membrane. The membranes were incubated overnight with primary antibodies: *TCL1A*, *TLR2*, *TLR7*, *TLR9*, *TLR10*, *MYD88*, *UNC93B1* (Novus Biologicals, Littleton, CO), and *ACTB* at a 1:500 dilution at 4°C. The washed membranes were then incubated with goat anti-rabbit or anti-mouse secondary antibody (Santa Cruz Biotechnology, Dallas, TX) at a 1:20,000 dilution. The washed membranes were subsequently incubated in Pierce ECL Western blotting substrate (Thermo Scientific, Madison, WI) and were visualized using Geldoc (Bio-Rad Laboratories, Hercules, CA).

**NF- $\kappa$ B Reporter Assay.** Lymphoblastoid cell lines with known *TCL1A* SNP genotypes were transfected by electroporation with 2  $\mu$ g of the pGL4.32[luc2P/NF- $\kappa$ B-RE.Hydro] vector (Promega, Madison, WI). A Renilla construct, pRL-TK (Promega), was used to determine transfection efficiency. The cells were plated in RPMI medium for 24 hours to allow them to recover from electroporation. Cells were then treated with 0.1 nM E<sub>2</sub> for 24 hours, followed by  $10^{-7}$   $\mu$ M 4-OH-TAM or ICI for an additional 24 hours. In some experiments, cells were treated with


MYD88 (100  $\mu$ M) for 24 hours before  $E_2$  treatment. Luciferase assays were performed with the Dual-Luciferase Reporter assay system (Promega). Relative NF- $\kappa$ B/luciferase activities were normalized to Renilla luciferase activities.

**Coimmunoprecipitation of TCL1A and MYD88.** LCLs ( $1 \times 10^7$ ) were resuspended in 650  $\mu$ l of immunoprecipitation (IP) lysis buffer containing 2.2  $\mu$ l of protease inhibitor cocktail (Qiagen, Valencia, CA) and incubated on ice for 30 minutes. Cells were then centrifuged at 12,000g at 4°C for 15 minutes. Supernatant was collected. Protein A agarose (Thermo Scientific) was prepared and washed three times with IP lysis buffer. A precleaning step was performed to clean the background. Cell lysates containing protein A agarose beads were rotated at 4°C for an hour. After centrifugation, supernatant was collected. At this point, input (50  $\mu$ l) was collected and stored at -80°C. Anti-TCL1A (1:50) antibodies (Cell Signaling Technology, Danvers, MA) were used to perform IP. IgG (Cell Signaling Technology), used as negative control. Specifically, IP samples containing protein A agarose beads were rotated at 4°C overnight. Immunoprecipitates were washed three times with ice cold lysis buffer, and proteins were eluted with 50  $\mu$ l of 1 $\times$  Laemmli loading buffer. Proteins were separated on 4%–12% SDS-PAGE gels and transferred onto polyvinylidene fluoride membranes. After blocking, membranes were incubated with primary antibodies against TCL1A at 4°C overnight. The washed membrane was then incubated with secondary antibody (1:15,000 dilution) for an hour at room temperature. The membrane was visualized using super-signal ECL substrate (Thermo Scientific).

**Statistics.** Data were analyzed using GraphPad Prism Software (San Diego, CA) and an R package. Data are presented as mean  $\pm$  S.E.M. unless otherwise stated. Gene expression and NF- $\kappa$ B activities were analyzed using multivariate analysis of variance, followed by planned comparisons with appropriate post hoc tests. We also conducted *t* tests for all meaningful combinations (effect of treatments on WT and variant, effect of genotype on treatments vs. vehicle). To correct for the inflated type 1 error that results from multiple comparisons, we have obtained a false discovery rate for each comparison using the R package (the default “p.adjust” function with “BH” method).

## Results

**Correlation between the Expression of TCL1A and that of Toll-Like Receptors.** Excessive production of cytokines and chemokines mediated by toll-like receptor (TLR) pathways is often associated with an inflammatory response mediated through NF- $\kappa$ B signaling (Zeytun et al., 2010). The human TLR family consists of 10 isoforms (TLR1–TLR10). TLRs are synthesized in the endoplasmic reticulum (Kawasaki and Kawai, 2014). Most TLRs are located on the cell membrane, but TLR3, TLR7, TLR8, and TLR9 are located on the membranes of intracellular compartments, such as the endoplasmic reticulum, endosomes, and lysosomes (Fig. 2) (Joosten et al., 2016). The myeloid differentiation primary response gene 88 (MYD88) encodes one of the functional adapter molecules that has not been shown to interact with all TLRs except TLR3. MYD88 recruits interleukin-1R-activating kinase (IRAK) 1, IRAK2, IRAK4, and tumor necrosis factor receptor-associated factor 6, leading ultimately to NF- $\kappa$ B activation, proinflammatory cytokine secretion, and an inflammatory response (Fig. 2). In the present study, in an attempt to extend our observations with regard to *TCL1A* SNPs and estrogen-dependent expression of immune mediators, we set out to determine whether TLR expression might also be regulated by the *TCL1A* SNPs observed in our GWAS in an estrogen-dependent fashion and, if so, whether these observations might cast light on the role of *TCL1A* SNPs in NF- $\kappa$ B activation.



**Fig. 2.** TLR intracellular localization and signaling. TLR3, TLR7, TLR8, and TLR9 are located on the membranes of intracellular compartments, such as the endoplasmic reticulum and endosomes. The myeloid differentiation primary response gene 88 (MYD88) is one of the functional adapter molecules that have been reported to interact with all TLRs except TLR3. MYD88 recruits IL-IRAK1, IRAK2, IRAK4, and tumor necrosis factor-receptor associated factor 6 (TRAF6), leading ultimately to NF- $\kappa$ B activation and proinflammatory cytokine secretion and an inflammatory response. UNC93B1 is a multitransmembrane-domain-containing protein and plays a critical role in trafficking TLR7 and TLR9 from the endoplasmic reticulum to endosomes, where TLR7 and TLR9 transmit signals via MYD88/TRIF-dependent pathways (Kim et al., 2008).

We once again turned to the Human Variation Panel of LCLs, a cell line model system that has repeatedly demonstrated its power to generate pharmacogenomic hypotheses and to test hypotheses arising from clinical GWA studies (Li et al., 2008; Ingle et al., 2010, 2013, 2016; Niu et al., 2010, 2016; Liu et al., 2013, 2014; Ho et al., 2016). This model system consists of LCLs from 300 healthy subjects for whom we have generated seven million SNPs after imputation, as well as microarray data for basal levels of gene expression for each cell line. We began by asking whether TLR expression might be associated with *TCL1A* gene expression. Basal levels of expression, in the absence of  $E_2$ , showed a significant correlation in these 300 cell lines of *TCL1A* expression with that of both TLR7 and TLR9, with *P* values of 6.43E-15 and 1.05E-09, respectively (Table 1). We also examined the correlation between the expression of MYD88, a critical adapter protein for inflammation signaling pathways downstream of TLRs (Takeuchi and Akira, 2010) and that of *Unc93 Homolog B1* (UNC93B1), a molecule that plays a critical role in trafficking TLR7 and TLR9 from the endoplasmic reticulum to endosomes (Fig. 2) (Sasai and Iwasaki, 2011; Lee and Barton, 2014). Very significant correlations were also observed between basal expression levels of TLR9 and TLR7 with the expression of MYD88, with *P* values 1.71E-18 and 7.55E-09, respectively (Table 2). We also found that expression levels of MYD88 and UNC93B1 were themselves significantly correlated ( $r = 0.41$ ,  $P = 7.66E-13$ ) (Table 2); however, in the absence of  $E_2$ , there was no significant SNP-dependent difference in the expression of the genes listed in Table 1 in these 300 LCLs (data not shown). Therefore, we also asked whether the expression of these genes might be influenced by

TABLE 1

Correlations of *TCL1A* mRNA expression and those of toll-like receptors and *MYD88* in the Human Variation Panel of 300 LCLs

Gene	Gene	<i>R</i>	<i>P</i> Value
<i>TCL1A</i>	<i>TLR1</i>	-0.089	1.36E-01
<i>TCL1A</i>	<i>TLR2</i>	0.276	2.54E-06
<i>TCL1A</i>	<i>TLR3</i>	-0.051	3.98E-01
<i>TCL1A</i>	<i>TLR4</i>	0.158	8.02E-03
<i>TCL1A</i>	<i>TLR5</i>	0.027	6.47E-01
<i>TCL1A</i>	<i>TLR6</i>	-0.073	2.21E-01
<i>TCL1A</i>	<i>TLR7</i>	0.443	6.43E-15
<i>TCL1A</i>	<i>TLR8</i>	0.052	3.89E-01
<i>TCL1A</i>	<i>TLR9</i>	0.354	1.05E-09
<i>TCL1A</i>	<i>TLR10</i>	-0.200	7.29E-04
<i>TCL1A</i>	<i>MYD88</i>	0.273	3.50E-06

Statistical significance was considered  $P < 1.8E-07$ .

the *TCL1A* SNPs in the presence of E<sub>2</sub>, as we had observed for proinflammatory cytokines, chemokines, and their receptors (Fig. 1, D–G) (Ho et al., 2016).

**TLR Expression Can Be Regulated by *TCL1A* SNPs in an Estrogen-Dependent Fashion.** The first set of experiments used eight LCLs that were homozygous for WT and eight LCLs homozygous for variant genotypes for the *TCL1A* SNPs that we had shown to be associated with AI-induced musculoskeletal adverse events (Ingle et al., 2010) and that we had also shown to be associated with alterations in the expression of a series of cytokines and chemokines in a SNP- and estrogen-dependent fashion (Fig. 1, D–G) (Ho et al.). All the cell lines used were homozygous for either WT or variant genotype for all three *TCL1A* SNPs, as shown graphically in Fig. 1A. *TCL1A* and all the immune mediators listed in Table 1 are highly expressed in LCLs—which are Epstein-Barr virus-transformed B cells—which made the functional genomic studies described subsequently possible.

The initial experiment represented an attempt to determine whether the expression of TLRs and *MYD88* could be modulated by *TCL1A* in a SNP-dependent fashion in response to E<sub>2</sub> treatment, and to determine whether the direction of that change in expression could be reversed in response to 4-OH-TAM treatment—implying that the expression might be regulated by *TCL1A* genotype. Concentrations of E<sub>2</sub> and 4-OH-TAM used to perform these experiments were those that were found to be optimal in previous studies of *TCL1A* induction by E<sub>2</sub> and the reversal of that effect by 4-OH-TAM (Liu et al., 2012; Ho et al., 2016). *TCL1A* expression was induced by 0.1 nM E<sub>2</sub> treatment, but only in cells homozygous for variant genotypes for the *TCL1A* SNPs (Fig. 3A)—confirming our previous results (Liu et al., 2012). This genotype-dependent gene expression pattern was reversed, however, when 4-OH-TAM ( $10^{-7}$   $\mu$ M) was added to E<sub>2</sub>—also confirming our previous results (Fig. 3A) (Ho et al., 2016). In parallel, the expression of *TLR2*, *TLR7*, *TLR9*, and *TLR10* were upregulated (~2- to 3-fold) in the presence of E<sub>2</sub>, but only in cells homozygous for variant genotypes for the *TCL1A* SNPs. In addition, as anticipated, in the presence of 4-OH-TAM, expression levels for these four TLRs increased significantly in cells homozygous for WT genotypes for the *TCL1A* SNPs (Fig. 3, B–E). Furthermore, other TLRs (*TLR1*, *TLR3*, *TLR4*, *TLR5*, *TLR6*, and *TLR8*) did not display alternation in their expression in a *TCL1A* SNP-dependent fashion in the presence of E<sub>2</sub> (data not shown). In a similar fashion, we also observed that the expression of *MYD88* and *UNC93B1* were significantly upregulated in parallel with *TCL1A* in a SNP-estrogen-dependent

TABLE 2

Correlations of *MYD88* mRNA expression with those of toll-like receptors and *UNC93B1* in the Human Variation Panel of 300 LCLs


Gene	Gene	<i>R</i>	<i>P</i> Value
<i>MYD88</i>	<i>TLR1</i>	0.19	1.36E-03
<i>MYD88</i>	<i>TLR2</i>	0.227	1.26E-04
<i>MYD88</i>	<i>TLR3</i>	0.061	3.10E-01
<i>MYD88</i>	<i>TLR4</i>	0.244	3.50E-05
<i>MYD88</i>	<i>TLR5</i>	0.057	3.42E-01
<i>MYD88</i>	<i>TLR6</i>	0.02	7.34E-01
<i>MYD88</i>	<i>TLR7</i>	0.491	1.71E-18
<i>MYD88</i>	<i>TLR8</i>	-0.097	1.03E-01
<i>MYD88</i>	<i>TLR9</i>	0.336	7.55E-09
<i>MYD88</i>	<i>TLR10</i>	0.145	1.50E-02
<i>MYD88</i>	<i>UNC93B1</i>	0.41	7.66E-13

Statistical significance was considered  $P < 1.8E-07$ .

fashion (Fig. 3, F–G). In addition, expression of those immune genes was significantly changed at the protein level as determined by Western blot analysis (Fig. 3H). It should be pointed out once again that these SNPs were in the *TCL1A* gene, as shown in Fig. 1A, not in genes encoding the immune mediators, that is, not in the *TLR2*, *TLR7*, *TLR9*, *TLR10*, *MYD88*, and *UNC93B1* genes (Fig. 3).

**Knockdown of *TCL1A* Resulted in Decreased Expression of TLRs.** We next determined whether *TCL1A* itself was involved in the differences in expression for *TLR2*, *TLR7*, *TLR9*, *TLR10*, *MYD88*, and *UNC93B1* shown in Fig. 3 using cell lines with differing genotypes for the *TCL1A* SNPs. Specifically, siRNA knockdown studies were performed using four independent siRNAs, as well as one pooled siRNA (Dharmacon Chicago, IL), and the results for all of these approaches were consistent. Knockdown of *TCL1A* to 25% of its baseline level resulted in significant downregulation of the expression of *TLR2*, *TLR7*, *TLR9*, *TLR10*, *MYD88*, and *UNC93B1*. We observed no SNP-dependent effect when *TCL1A* was knocked down (Fig. 4, A and B). In a similar fashion, knockdown of *MYD88* resulted in significantly decreased *TLR2*, *TLR7*, *TLR9*, and *TLR10* expression in LCLs, but *TCL1A* expression did not change (Fig. 4, D and E). Finally, protein expression was altered in parallel with the changes seen for mRNA after *TCL1A* or *MYD88* knockdown (Fig. 4, C and F). These results indicated that the expression of *TLR2*, *TLR7*, *TLR9*, and *TLR10* could be modulated by *TCL1A* and *MYD88*. It should be pointed out that knockdown of *MYD88* did not alter the expression of *TCL1A*, suggesting that *TCL1A* is upstream of *MYD88* in this signaling pathway. At this point, we had determined that *TCL1A* can regulate the expression of TLRs as well as that of *MYD88*, an important adaptor molecule that can trigger downstream signaling, including signaling through the NF- $\kappa$ B pathway.


***TCL1A* SNP and SERM-Mediated *MYD88*-Dependent NF- $\kappa$ B Activation.** We had previously reported that changes in *TCL1A* expression after ER blockade using fulvestrant (ICI) could influence NF- $\kappa$ B transcriptional activity in a striking SNP genotype-dependent fashion (Liu et al., 2012). Specifically, we treated the same cells that were used to generate the results shown in Fig. 3, LCLs with differing genotypes for the *TCL1A* genotypes, with ICI or 4-OH-TAM. Cells homozygous for the variant genotype for the *TCL1A* SNPs increased NF- $\kappa$ B transcriptional activity approximately 3-fold in the presence of E<sub>2</sub> plus either ICI or 4-OH-TAM (Fig. 5, A and B). *MYD88* is the adaptor protein for all TLRs except *TLR3* and is capable of activating the NF- $\kappa$ B signaling pathway and of inducing the



**Fig. 3.** SNP- and estrogen-dependent mRNA expression of *TCL1A* (A), *TLR2* (B), *TLR7* (C), *TLR9* (D), *TLR10* (E), *MYD88* (F), and *UNC93B1* (G) in LCLs (H). Western blot analysis was performed for *TCL1A*, *TLR2*, *TLR7*, *TLR9*, *TLR10*, *MYD88*, *UNC93B1*, and *ACTB* in LCLs with known *TCL1A* SNP genotypes. The cells were treated with 0.1 nM E<sub>2</sub> or with 0.1 nM E<sub>2</sub> plus 10<sup>-7</sup> μM 4-hydroxytamoxifen (4OH-TAM) for an additional 24 hours. Eight cell lines homozygous for the variant (V) genotypes for all three of the *TCL1A* SNPs and eight cell lines homozygous for WT genotypes were used in these experiments. \*\*\**P* < 0.0001.

production of inflammatory cytokines (Broad et al., 2007; Kawai and Akira, 2007). We had observed that MYD88 expression could be regulated by *TCL1A* in a SNP- and estrogen-dependent fashion (Fig. 3F) and that it is downstream of *TCL1A* (Fig. 4, A and B). The next series of experiments was designed to study the possible contribution of MYD88, a known adaptor molecule for TLR-mediated NF-κB signaling, to *TCL1A* SNP and SERM-dependent NF-κB activation. We found that, after knockdown of

MYD88 to ~30% of its baseline level, *TCL1A* SNP-dependent NF-κB transcriptional activation in the presence of either 4-OH-TAM or ICI was lost (Fig. 5C). In addition, when MYD88 was silenced by the application of MYD88 inhibitory peptide, the effects of the *TCL1A* SNP-dependent and ER blocker-dependent NF-κB activation were also lost, as anticipated (Fig. 5D). We also determined by coimmunoprecipitation that *TCL1A* could interact directly with MYD88 (Fig. 5E). These results further



**Fig. 4.** TCL1A could modulate TLR2, TLR7, TLR9, TLR10, MYD88, and UNC93B1 expression in LCLs. Relative mRNA expression (A and B) of TCL1A, TLR2, TLR7, TLR9, TLR10, MYD88, and UNC93B1 after knockdown of TCL1A in LCLs with known *TCL1A* SNP genotypes using pooled siRNA. Eight cell lines of each genotype were used in these experiments. \* $P < 0.05$ . Student's *t* test was performed to compare gene expression in LCLs with differing *TCL1A* SNP genotypes before and after gene knockdown. \* $P$  value  $\leq 0.05$  was considered statistically significant. All values are mean  $\pm$  S.E.M for three separate independent assays. Protein expression was determined by Western blot analysis (C). Relative mRNA expression (D and E) of TCL1A, TLR2, TLR7, TLR9, TLR10, MYD88, and UNC93B1 after knockdown of MYD88 in LCLs with known *TCL1A* SNP genotypes using pooled siRNA. (F) Western blot analysis was performed for TCL1A, TLR2, TLR7, TLR9, TLR10, MYD88, and UNC93B1 after knockdown of MYD88.


confirmed that *TCL1A* SNP and estrogen-dependent NF- $\kappa$ B activation may occur, in part, through a TLR-MYD88-dependent pathway. They also greatly extended our original observations and served to highlight a novel pharmacogenomic mechanism by which *TCL1A* can influence and modulate the immune response and inflammation.

## Discussion

We previously reported that SNPs 3' of the *TCL1A* gene that we identified during a GWAS for musculoskeletal adverse events that occur during aromatase inhibitor therapy of breast cancer could have significant effects on the expression of a series of genes encoding inflammatory mediators in an SNP- and estrogen-dependent fashion. Even more striking, those effects could be reversed in a *TCL1A* SNP-dependent fashion by exposure to 4-OH-TAM or ICI (Ho et al., 2016). We also reported that variant *TCL1A* SNP genotypes were associated

with increased NF- $\kappa$ B transcriptional activity after ER blockade, but that was not true for the WT genotype (Liu et al., 2012). NF- $\kappa$ B activation is known to play an important role in chronic inflammation (Hayden et al., 2006). As a result, there is great interest in developing specific molecular inhibitors of NF- $\kappa$ B activation. Therefore, the present study represents a step toward a better understanding of molecular and genetic mechanisms responsible for variation in NF- $\kappa$ B activation—understanding that could potentially have implications for the management of NF- $\kappa$ B-related inflammation.

In the present study, we pursued mechanisms associated with the *TCL1A* SNP-mediated, SERM-dependent activation of NF- $\kappa$ B. It has been reported that TCL1A can function as a transcriptional regulator, which interacts with many transcription factors, including AP1 and cAMP response element binding/p300 and, as a result, that it can influence NF- $\kappa$ B activity in B-cell chronic lymphocytic leukemia (Pekarsky et al., 2008). Since TLR signaling, signaling that occurs—in



**Fig. 5.** *TCL1A* SNP- and estrogen-dependent NF-κB activation as determined by NF-κB reporter assays could be altered by the knockdown or inhibition of MYD88. ER blockade by fulvestrant (ICI) or 4-hydroxytamoxifen (4OH) treatment resulted in *TCL1A* SNP-dependent NF-κB activation (A and B). *TCL1A* SNP and estrogen-dependent NF-κB activation could be blocked by MYD88 siRNA knockdown (C) or by exposure to MYD88 inhibitory peptide (100 μM) (D). Specifically, LCLs were cotransfected with an NF-κB reporter construct and siRNA (MYD88 or control siRNA); 24 hours after transfection, cells were treated with either vehicle or 0.1 nM E<sub>2</sub> for 24 hours, followed by 10<sup>-7</sup> μM 4OH or ICI for an additional 24 hours. In some experiments, cells were exposed to MYD88 inhibitory peptide for 24 hours before E<sub>2</sub> treatment. Luciferase activity was measured 72 hours after transfection. The firefly luciferase activity derived from the NF-κB responsive reporter was normalized by the use of Renilla luciferase activity as a control to correct for possible variation in transfection efficiency. All experiments were repeated three times in triplicate. \*\**P* < 0.001. (E) Coimmunoprecipitation was used to determine whether *TCL1A* protein could interact with MYD88 in LCLs. Whole-cell lysates from 1 × 10<sup>7</sup> LCLs were immunoprecipitated with anti-*TCL1A* (1:50) antibodies or anti-IgG antibodies. Whole-cell lysate (input, left panel) and immunoprecipitated samples (middle and right panels) were immunoblotted and probed with antibodies against *TCL1A* and MYD88.

part—through MYD88, can influence NF-κB activity (Broad et al., 2007), MYD88 has been suggested as a potential therapeutic target for regulating the immune response (Olson et al., 2015). When *TCL1A* was knocked down in our cell line model system, MYD88 was significantly down-regulated and, in parallel, expression levels of TLR2, TLR7, TLR9, and TLR10 were also decreased significantly (Fig. 4, C and D). Several TLRs can interact with MYD88 and NF-κB, as determined by STRING protein interaction network analysis (Szklarczyk et al., 2015). We also observed by IP studies that *TCL1A* protein could interact with MYD88 (Fig. 5E). We have reported previously that ER blockade by fulvestrant (ICI) can result in the activation of NF-κB transcriptional activity, but only in cells with variant genotypes for the *TCL1A* SNPs (3) (Fig. 5A). In the present study, we extended that finding by observing a similar response after exposure to 4-OH-TAM (Fig. 5B). We also

demonstrated that the *TCL1A* SNP- and SERM-dependent activation of NF-κB through TLRs could be blocked by MYD88 siRNA or MYD88 inhibitory peptide (Fig. 5, C and D). Therefore, the present study has raised the possibility that TLR-MYD88-dependent NF-κB signaling could be involved in *TCL1A* SNP-dependent NF-κB activation and, as a result, in the regulation of inflammation and immune response. The fact that expression of a single gene, *TCL1A*, can influence signaling through this pathway represents an important advance in our understanding of the regulation of the pathway and suggests novel ways to influence signaling through this pathway.

Obviously, further studies will be required to determine the possible therapeutic implications of this series of observations. Specifically, we found that *TCL1A* expression was positively correlated with the basal expression of toll-like receptors TLR7 and TLR9 in 300 LCLs (*r* = 0.443, *P* = 6.43E-15, and



$r = 0.354$ ,  $P = 1.05E-09$ , respectively), as listed in Table 1. Both TLR7 and TLR9 are therapeutic targets for the treatment of lupus erythematosus, an autoimmune disorder for which more than 90% of patients are women. We also found that TCL1A knockdown could influence the expression of TLR2 and TLR10, both of which can play a role in cytokine production, inflammation, and NF- $\kappa$ B activation via MYD88 (Joosten et al., 2016). Crosstalk between estrogens and the immune response is well documented (Seillet et al., 2012; Hughes and Choubey, 2014; Kovats, 2015). In the present study, we have demonstrated the differential expression of a series of genes encoding immune mediators in response to estrogen treatment that was dependent on *TCL1A* SNP genotypes. Even more striking, we showed that 4-OH-TAM, an active metabolite of tamoxifen, a Food and Drug Administration–approved treatment of breast cancer and for breast cancer chemopreventive that has been suggested for the treatment of millions of women at high risk for breast cancer (Maximov et al., 2013), can reverse the *TCL1A* SNP-dependent expression of those immune mediators—all of which play a critical role in inflammatory disease (Iwakura et al., 2011; Bartlett and Million, 2015).

We should emphasize that the LCL model system used in our studies has both strengths and weaknesses, but it has repeatedly demonstrated that it can be a powerful tool for both generating and testing genomic and pharmacogenomics hypotheses (Li et al., 2008, 2012, 2013, 2014; Ingle et al., 2010, 2013; Niu et al., 2010; Wheeler and Dolan, 2012). The availability of comprehensive genotype and gene expression data for this model system makes it possible to study the functional implications of genetic variants found to be associated with clinical phenotypes, as demonstrated by the present studies. We have previously used this model system successfully to demonstrate novel SNP- and estrogen-dependent mechanisms underlying variation in the regulation of the expression of cytokines and chemokines by *TCL1A* (Liu et al., 2012). In the present study, we have extended those observations with regard to the transcriptional regulation of immune mediators (cytokines and chemokines) to include TLRs and NF- $\kappa$ B transcriptional activity. Obviously, these results will have to be verified using both additional cell-based systems and clinical samples, but the present results represent an important step in the process of providing functional and mechanistic explanations for the association of SNPs in the *TCL1A* gene with inflammation and the immune response.

In summary, the series of functional genomic studies described here confirm that *TCL1A* expression is  $E_2$  inducible in an SNP-dependent manner and have demonstrated that *TCL1A* can influence the downstream expression of a series of immune mediators, including TLR2, TLR7, TLR9, TLR10, and MYD88. Furthermore, inhibition of MYD88 resulted in the blockade of *TCL1A* SNP-dependent NF- $\kappa$ B activation, indicating that the TLR-MYD88–dependent NF- $\kappa$ B signaling pathway might contribute, at least in part, to the *TCL1A* SNP- and estrogen-dependent effects that we had observed. *TCL1A* SNP- and estrogen-dependent variation in NF- $\kappa$ B transcriptional activity and the transcriptional regulation of other immune mediators suggests that this pathway may play a role in the complex interactions that are known to exist between the endocrine and immune systems. Of equal importance is the fact that SERMs are able to modulate this pathway in a striking fashion—opening the way for the pharmacologic regulation of this pathway in clinical setting.

#### Authorship Contributions

*Participated in research design:* Ho, Ingle, Bongartz, Goss, Shepherd, Mushiroda, Kubo, Wang, Weinshilboum.

*Conducted experiments:* Ho.

*Performed data analysis:* Ho, Ingle, Bongartz, Wang, Weinshilboum.

*Wrote or contributed to the writing of the manuscript:* Ho, Ingle, Bongartz, Kalari, Goss, Shepherd, Mushiroda, Kubo, Wang, Weinshilboum.

#### References

- Bartlett HS and Million RP (2015) Targeting the IL-17(T/H)17 pathway. *Nat Rev Drug Discov* **14**:11–12.
- Broad A, Kirby JA, and Jones DEJ; Applied Immunology and Transplantation Research Group (2007) Toll-like receptor interactions: tolerance of MyD88-dependent cytokines but enhancement of MyD88-independent interferon- $\beta$  production. *Immunology* **120**:103–111.
- Hayden MS, West AP, and Ghosh S (2006) NF- $\kappa$ B and the immune response. *Oncogene* **25**:6758–6780.
- Ho M-F, Bongartz T, Liu M, Kalari KR, Goss PE, Shepherd LE, Goetz MP, Kubo M, Ingle JN, Wang L, et al. (2016) Estrogen, SNP-dependent chemokine expression and selective estrogen receptor modulator regulation. *Mol Endocrinol* **30**:382–398.
- Hughes GC and Choubey D (2014) Modulation of autoimmune rheumatic diseases by oestrogen and progesterone. *Nat Rev Rheumatol* **10**:740–751.
- Ingle JN, Liu M, Wickerham DL, Schaid DJ, Wang L, Mushiroda T, Kubo M, Costantino JP, Vogel VG, Paik S, et al. (2013) Selective estrogen receptor modulators and pharmacogenomic variation in ZNF423 regulation of BRCA1 expression: individualized breast cancer prevention. *Cancer Discov* **3**:812–825.
- Ingle JN, Schaid DJ, Goss PE, Liu M, Mushiroda T, Chapman J-AW, Kubo M, Jenkins GD, Batzler A, Shepherd L, et al. (2010) Genome-wide associations and functional genomic studies of musculoskeletal adverse events in women receiving aromatase inhibitors. *J Clin Oncol* **28**:4674–4682.
- Ingle JN, Xie F, Ellis MJ, Goss PE, Shepherd LE, Chapman JW, Chen BE, Kubo M, Furukawa Y, Momozawa Y, et al. (2016) Genetic polymorphisms in the long non-coding RNA MIR2052HG offer a pharmacogenomic basis for the response of breast cancer patients to aromatase inhibitor therapy. *Cancer Res* **76**:7012–7023.
- Iwakura Y, Ishigame H, Saijo S, and Nakae S (2011) Functional specialization of interleukin-17 family members. *Immunity* **34**:149–162.
- Joosten LAB, Abdollahi-Roodsaz S, Dinarello CA, O'Neill L, and Netea MG (2016) Toll-like receptors and chronic inflammation in rheumatic diseases: new developments. *Nat Rev Rheumatol* **12**:344–357.
- Kawai T and Akira S (2007) Signaling to NF- $\kappa$ B by Toll-like receptors. *Trends Mol Med* **13**:460–469.
- Kawasaki T and Kawai T (2014) Toll-like receptor signaling pathways. *Front Immunol* **5**:461.
- Kim Y-M, Brinkmann MM, Paquet M-E, and Ploegh HL (2008) UNC93B1 delivers nucleotide-sensing toll-like receptors to endolysosomes. *Nature* **452**:234–238.
- Kovats S (2015) Estrogen receptors regulate innate immune cells and signaling pathways. *Cell Immunol* **294**:63–69.
- Lee BL and Barton GM (2014) Trafficking of endosomal Toll-like receptors. *Trends Cell Biol* **24**:360–369.
- Li L, Fridley B, Kalari K, Jenkins G, Batzler A, Safgren S, Hildebrandt M, Ames M, Schaid D, and Wang L (2008) Gemcitabine and cytosine arabinoside cytotoxicity: association with lymphoblastoid cell expression. *Cancer Res* **68**:7050–7058.
- Liu M, Goss PE, Ingle JN, Kubo M, Furukawa Y, Batzler A, Jenkins GD, Carlson EE, Nakamura Y, Schaid DJ, et al. (2014) Aromatase inhibitor-associated bone fractures: a case-cohort GWAS and functional genomics. *Mol Endocrinol* **28**:1740–1751.
- Liu M, Ingle JN, Fridley BL, Buzdar AU, Robson ME, Kubo M, Wang L, Batzler A, Jenkins GD, Pietrzak TL, et al. (2013) TSPYL5 SNPs: association with plasma estradiol concentrations and aromatase expression. *Mol Endocrinol* **27**:657–670.
- Liu M, Wang L, Bongartz T, Hawse JR, Markovic SN, Schaid DJ, Mushiroda T, Kubo M, Nakamura Y, Kamatani N, et al. (2012) Aromatase inhibitors, estrogens and musculoskeletal pain: estrogen-dependent T-cell leukemia 1A (*TCL1A*) gene-mediated regulation of cytokine expression. *Breast Cancer Res* **14**:R41.
- Maximov PY, Lee TM, and Jordan VC (2013) The discovery and development of selective estrogen receptor modulators (SERMs) for clinical practice. *Curr Clin Pharmacol* **8**:135–155.
- McCool KW and Miyamoto S (2012) DNA damage-dependent NF- $\kappa$ B activation: NEMO turns nuclear signaling inside out. *Immunol Rev* **246**:311–326.
- Niu N, Liu T, Cairns J, Ly RC, Tan X, Deng M, Fridley BL, Kalari KR, Abo RP, Jenkins G, et al. (2016) Metformin pharmacogenomics: a genome-wide association study to identify genetic and epigenetic biomarkers involved in metformin anti-cancer response using human lymphoblastoid cell lines. *Hum Mol Genet* **25**:4819–4834.
- Niu N, Qin Y, Fridley BL, Hou J, Kalari KR, Zhu M, Wu T-Y, Jenkins GD, Batzler A, and Wang L (2010) Radiation pharmacogenomics: a genome-wide association approach to identify radiation response biomarkers using human lymphoblastoid cell lines. *Genome Res* **20**:1482–1492.
- Olson MA, Lee MS, Kissner TL, Alam S, Waugh DS, and Saikh KU (2015) Discovery of small molecule inhibitors of MyD88-dependent signaling pathways using a computational screen. *Sci Rep* **5**:14246.
- Pekarsky Y, Palamarchuk A, Maximov V, Efanov A, Nazaryan N, Santanam U, Rassenti L, Kipps T, and Croce CM (2008) Tc1 functions as a transcriptional regulator and is directly involved in the pathogenesis of CLL. *Proc Natl Acad Sci USA* **105**:19643–19648.
- Sasai M and Iwasaki A (2011) Love triangle between Unc93B1, TLR7, and TLR9 prevents fatal attraction. *Immunity* **35**:3–5.

- Seillet C, Laffont S, Trémollières F, Rouquié N, Ribot C, Arnal J-F, Douin-Echinard V, Gourdy P, and Guéry J-C (2012) The TLR-mediated response of plasmacytoid dendritic cells is positively regulated by estradiol in vivo through cell-intrinsic estrogen receptor  $\alpha$  signaling. *Blood* **119**:454–464.
- Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, Simonovic M, Roth A, Santos A, Tsafou KP, et al. (2015) STRING v10: protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res* **43**: D447–D452.
- Tak PP and Firestein GS (2001) NF-kappaB: a key role in inflammatory diseases. *J Clin Invest* **107**:7–11.
- Takeda K, Harada Y, Watanabe R, Inutake Y, Ogawa S, Onuki K, Kagaya S, Tanabe K, Kishimoto H, and Abe R (2008) CD28 stimulation triggers NF-kappaB activation through the CARMA1-PKCtheta-Grb2/Gads axis. *Int Immunol* **20**:1507–1515.
- Takeuchi O and Akira S (2010) Pattern recognition receptors and inflammation. *Cell* **140**:805–820.
- Wheeler HE and Dolan ME (2012) Lymphoblastoid cell lines in pharmacogenomic discovery and clinical translation. *Pharmacogenomics* **13**:55–70.
- Zeytun A, Chaudhary A, Pardington P, Cary R, and Gupta G (2010) Induction of cytokines and chemokines by toll-like receptor signaling: strategies for control of inflammation. *Crit Rev Immunol* **30**:53–67.

---

**Address correspondence to:** Richard Weinshilboum, M.D., Mayo Clinic 200 First Street SW, Rochester, MN 55905. E-mail: weinshilboum.richard@mayo.edu

---